

NO GW IS EMITTED BY B PSR1913+16

ANGELO LOINGER

ABSTRACT. In the *exact* (non-linear) formulation of general relativity (GR) *no* motion of bodies can give origin to gravitational waves (GW's) – as it has been proved. Accordingly, the measured rate of change of the orbital period of binary pulsar B PSR1913+16 must have *other* causes, different from the emission of GW's; maybe the viscous losses of the unseen pulsar companion, if it were e.g. a helium star.

Summary. - The paper is structured as follows. Sect.1. contains some passages of a recent report by Weisberg and Taylor (see [1*g*]) regarding thirty years of observations of binary radiopulsar B PSR1913+16. Sect.2.: a straightforward criticism of the relativistic approach which is employed in papers cited in [1]. Sect.2*bis*.: a possible alternative explanation of the shrinkage of the orbit of the above radiopulsar. Sect.3.: the linear approximation of GR is *inadequate* to give an existence theorem of physical GW's. Sect.4.: *system B PSR1913+16 cannot emit GW's*. Sect.5.: the analogy between Maxwell-Lorentz e.m. theory and the linearized version of GR is a *false* analogy. Sect.6.: erroneousess of a surmise concerning the behaviour of B PSR1913+16.

1. - Nobody has ever found a ***direct*** experimental proof of the *real* existence of the gravitational waves (GW's). According to some authors [1], an ***indirect*** experimental evidence could be given by the time decrease of the orbital period P_b of the binary pulsar B PSR1913+16 [1*bis*].

The abstract of paper [1*g*]) runs as follows: “We describe results derived from thirty years of observations of PSR B1913+16. Together with the Keplerian orbital parameters, measurements of the relativistic periastron advance and a combination of gravitational redshift and time dilation yield the stellar masses with high accuracy. The measured rate of change of orbital period agrees with that expected from the emission of gravitational radiation, according to general relativity, to within about 0.2 percent. Systematic effects depending on the pulsar distance and on poorly known galactic constants now dominate the error budget, so tighter bounds will be difficult to obtain. [...]”. And in sect.3.1 of the same paper [1*g*]) the authors claim that: “According to general relativity, a binary star system should emit

To be published on *Spacetime & Substance*.
email: angelo.loinger@mi.infn.it
Dipartimento di Fisica, Università di Milano, Via Celoria, 16 - 20133 Milano (Italy).

energy in the form of gravitational waves. The loss of orbital energy results in shrinkage of the orbit, which is most easily observed as a decrease in orbital period. Peters and Mathews (1963) (see [2]) showed that in general relativity the rate of period decrease is given by

$$(1) \quad \dot{P}_{b,GR} = -\frac{192\pi G^{5/3}}{5c^5} \left(\frac{P_b}{2\pi}\right)^{-5/3} (1-e^2)^{-7/2} \times \left(1 + \frac{73e^2}{24} + \frac{37e^4}{96}\right) m_p m_c (m_p + m_c)^{-1/3}."$$

Here: G is the gravitational constant; c the speed of light *in vacuo*; e the orbital eccentricity ($e = 0.6171338(4)$); m_p the mass of the pulsar ($m_p = 1.4414 \pm 0.0002$ solar masses), m_c the mass of the companion ($m_c = 1.3867 \pm 0.0002$ solar masses).

Then, Weisberg and Taylor [1g)] write: "Comparison of the measured $\dot{P}_b$ with the theoretical value requires a small correction, $\dot{P}_{b,Gal}$, for relative acceleration between the solar system and binary pulsar system, projected onto the line of sight (Damour and Taylor 1991) [*see*[1e)]]. This correction is applied to the measured $\dot{P}_b$ to form a "corrected value" $\dot{P}_{b,corrected} = \dot{P}_b - \dot{P}_{b,Gal}$. The correction term depends on several rather poorly known quantities, including the distance [$\approx 16,000$ light-years] and proper motion of the pulsar and the radius of the Sun's galactic orbit. The best currently available values yield $\dot{P}_{b,Gal} = -(0.0128 \pm 0.0050) \times 10^{-12}$ s/s, so that $\dot{P}_{b,corrected} = 2.4056 \pm 0.0051) \times 10^{-12}$ s/s. Hence

$$(2) \quad \frac{\dot{P}_{b,corrected}}{\dot{P}_{b,GR}} = 1.0013 \pm 0.0021,$$

and we conclude that the measured orbital decay is consistent at the (0.13 ± 0.21) % level with the general relativistic prediction for the emission of gravitational radiation. [...]."

2. - The good agreement between the measured $\dot{P}_b$ and the computed $\dot{P}_b$ is suspect – as I have already emphasized [3] –, because the relativistic perturbative approximation, of which eq.(1) is a consequence, is quite unreliable from the point of view of the **exact** (non-linear) formulation of GR, as it was pointed out by several relativists [3bis].

Further, I remark that in GR the hypothetical GW's do not have a *true* energy. Therefore the *true* mechanical energy which is lost during the orbital motion should transform itself into the *pseudo* (i.e. false) energy of the hypothetical GW's: the energy balance would be violated. **Objection:** if we *suppose* that the *linearized* version of GR has an unconditioned, approximate validity – as the experimentalists, and some (simple) theoreticians [4], do – the physical existence of GW's seems a theoretical possibility, and it seems – by exploiting the analogy with Maxwell-Lorentz e.m. theory – that the *acceleration* of a body can generate GW's. **Answer:** the energy-momentum of such GW's has a tensor character only under Lorentz transformations of

co-ordinates, but *not* under general transformations. Now, this is contrary to the basic tenet of GR.

2bis. - The authors of papers [1] have *assumed* that *both* stars of the considered binary system are neutron stars, and thus act dynamically as *point* masses. But if the companion star were a helium star or a white dwarf, tides and viscous actions might mimic the relativistic (as the periastron advance) and pseudorelativistic effects. In particular, the viscous losses of the companion could give a time decrease of the pulsar revolution period of the same order of magnitude of that given by the hypothesized emission of gravitational radiation – as it is well known to many observational astrophysicists.

Finally, the empirical success of a theory – or of a given computation – is not an absolute guaranty for its conceptual adequacy. Consider for instance the Ptolemaic theory of cycles and epicycles, which explained rather well the planetary orbits (with the only exception of Mercury’s).

3. - It can be proved that the linear approximation of GR is quite *inadequate* to a proper study of the hypothetic GW’s, see [5], [6]. And if we continue the approximation beyond the linear stage (see [7], [8]), we find that the radiation terms of the gravitational field can be *destroyed* by convenient co-ordinate transformations: this proves that the GW’s are *only a product of a special choice of the reference system*, i.e. that they do not possess a *physical* reality (see further [9], [10], [11]): the undulatory solutions of Einstein field equations have a mere *formal* (*non-physical*) character.

4. - If the two stars of B PSR1913+16 are dynamically treated as two (gravitationally interacting) *point* masses [1], the *exact* formulation of GR tells us that their orbits are *geodetic* lines [9], i.e. their motions are “natural”, “free” motions, quite analogous to a rectilinear and uniform motion of a point charge in the customary Maxwell-Lorentz theory. Accordingly, no GW is sent forth by our stars! [10], [11].

In my paper [3 β] I have given another elegant proof of this fact, resting on a fundamental proposition by Hermann Weyl [12], according to which for any relative motion of two bodies it is always possible (in GR) to choose a co-ordinate system in which *both* bodies are *at rest*. (Remark that in GR the expression *at rest* must be defined precisely by specifying the *interested spacetime manifold*.)

Let us apply the above proposition to system B PSR1913+16, i.e. let us choose a co-ordinate frame for which both stars are at rest. Evidently, an observer Ω who “resides” in this frame does not record any emission of GW’s. Now, any observer Ω' – very far, in particular, from Ω –, for whom B PSR1913+16 is in motion, does not possess (in GR!) any physical privilege with respect to Ω . Accordingly, both observers, Ω and Ω' , do not register any GW sent forth by our binary system. (See Weyl [13] for the Riemann-Einstein manifold of two point masses at rest.)

5. - The *false* formal analogy between the e.m. Maxwell-Lorentz theory and the linearized version of GR is the responsible for the publication of

countless and senseless papers. In particular, it has generated the conviction that, in GR, the *acceleration* of a body must give origin to GW's; many people have forgotten that, in the *exact* (non-linear) formulation of GR, the concept "acceleration" does not possess an absolute character. (The above conviction was also extended to perturbative approximations of higher order.)

We observe finally that the exact theory does not admit any class of physically privileged reference frames for which, in particular, the undulatory character of a given gravitational field is an invariant property.

6. - A last remark. Some authors have conjectured that a coexistence of effects due to emission of GW's by B PSR1913+16 and to tides and viscous actions of the companion star could be possible.

Now, this is pure nonsense, because – as it can be proved – even motions that are *not* purely gravitational cannot generate GW's [14].

Πῦρ σοι προσοίσω.

(I will bring fire to thee.)

EURIPIDES, *Andromache*.

REFERENCES

- [1] : a) J.H. Taylor *et al.*, *Nature*, **277** (1979) 437; b) J.H. Taylor and J.M. Weisberg, *Astrophys. J.*, **253** (1982) 908; c) T. Damour and N. Deruelle, *Ann. Inst. Poincaré*, **43** (1985) 107; d) J.H. Taylor and J.M. Weisberg, *Astrophys. J.*, **345** (1989) 434; e) T. Damour and J.H. Taylor, *Astrophys. J.*, **366** (1991) 501; f) J.H. Taylor, *Revs. Mod. Phys.*, **66** (1994) 711; g) J.M. Weisberg and J.H. Taylor, *Binary Radio Pulsars*, *ASP Conference Series*, Vol. *TDB*, 2004; also on *arXiv:astro-ph/0407149 v1* (July 7th, 2004).
- [1bis] In his Nobel lecture – see *Revs. Mod. Phys.*, **66** (1994) 699 – R.A. Hulse does not discuss the question of the GW's, which would be emitted by B PSR1913+16.
- [2] P.C. Peters and J. Mathews, *Phys. Rev.*, **131** (1963) 435.
- [3] A. Loinger: α) *arXiv:astro-ph/9810137* (October 8th, 1998); β) *arXiv:astro-ph/0002267* (February 12th, 2000); γ) *Spacetime & Substance*, **3** (2002) 145; also on *arXiv:physics/0207013 v1* (July 2nd, 2002). For papers α) and β) see also: *On Black Holes and Gravitational Waves* (La Goliardica Pavese, Pavia) 2002, pp.41 and 72, respectively.
- [3bis] Cf., for instance, A. Rosenblum, *Phys. Rev. Letters*, **41** (1978) 1003.
- [4] Cf., e.g., the following papers: *i*) J. Hough, S. Rowan and B.S. Sathyaprakash, *arXiv:gr-qc/0501007 v1* (January 4th, 2005); *ii*) É. É. Flanagan and S. A. Hughes, *arXiv:gr-qc/0501041 v1* (January 12th, 2005).
- [5] H. Weyl, *Amer. J. Math.*, **66** (1944) 591.
- [6] A. Loinger, *Spacetime & Substance*, **5** (2004) 57; also on *arXiv:physics/0407134 v1* (July 27th, 2004).
- [7] L. Infeld and J. Plebanski, *Motion and relativity* (Pergamon Press, Oxford, etc.) 1960, in particular Chapt. VI.
- [8] A. Scheidegger, *Revs. Mod. Phys.*, **25** (1953) 451.
- [9] A. Loinger, *Nuovo Cimento B*, **115** (2000) 679; also on *arXiv:astro-ph/0003230* (March 16th, 2000).
- [10] For many exact proofs of the *physical* non-existence of the GW's, see A. Loinger, *arXiv:physics/0312149 v3* (11 February 2004) and the literature quoted there *sub* [6], in particular: A. Loinger, *Spacetime & Substance*, **3** (2002) 129 and **3** (2002) 145; Idem, *On Black Holes and Gravitational Waves* (La Goliardica Pavese, Pavia) 2002, Part II. See also: Idem, *arXiv: gr-qc, astro-ph, physics* (1998-2003).

- [11] The first theoretical proof of the *physical* non-existence of the GW's was given by T. Levi-Civita in 1917, see *Rend. Lincei*, **26** (1917) 381; an English version of this fundamental memoir in *arXiv:physics/9906004* (June 2nd, 1999).
- [12] H. Weyl, *Raum-Zeit-Materie*, Siebente Auflage (Springer-Verlag, Berlin, *etc.*) 1988, p.268.
- [13] See Weyl [12], pp.266-267.
- [14] See, in particular: A. Loinger, *On Black Holes and Gravitational Waves* (La Goliardica Pavese, Pavia) 2002, p.90 – also on *arXiv:physics/0106052* (June 17th, 2001); Idem, *Spacetime & Substance*, **3** (2002) 129 – also on *arXiv:physics/0202065 v1* (February 27th, 2002); Idem, *arXiv:physics/0412164 v1* (December 25th, 2004) – to be published in *Spacetime & Substance*.

★-----★